



# Adult Smoking - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Percent of adults that currently smoke and have smoked at least 100 cigarettes in lifetime, 2003-2009



## CONTEXT

**What It Is:** Adult smoking prevalence is the estimated percent of the adult population that currently smokes every day or “most days” and has smoked at least 100 cigarettes in their lifetime.

**Where It Comes From:** This measure was calculated by the National Center for Health Statistics using data from the Centers for Disease Control and Prevention’s Behavioral Risk Factor Surveillance System (BRFSS), a random-digit dial survey. BRFSS data are representative of the total non-institutionalized U.S. population ages 18 and older living in households with a land-line telephone. The estimates are based on seven years of data.


**Importance:** Each year approximately 443,000 premature deaths occur in the U.S. primarily due to smoking. Cigarette smoking is identified as a cause in multiple diseases including various cancers, cardiovascular disease, respiratory conditions, low birthweight, and other adverse health outcomes. Measuring the prevalence of tobacco use in the population can alert communities to potential adverse health outcomes and can be valuable for assessing the need for cessation programs or the effectiveness of existing programs.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Adult Obesity - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Percent of adults that report a body mass index (BMI) of at least 30 kg/m2, 2008



## CONTEXT

**What It Is:** The adult obesity measure represents the percent of the adult population (age 20 and older) that has a body mass index (BMI) greater than or equal to 30 kg/m2.

**Where It Comes From:** Estimates of obesity prevalence by county were calculated by the CDC’s National Center for Chronic Disease Prevention and Health Promotion, Division of Diabetes Translation, using multiple years of Behavioral Risk Factor Surveillance System (BRFSS) data. BRFSS data are representative of the total non-institutionalized U.S. population ages 18 and older living in households with a land-line telephone.


**Importance:** Obesity is often the end result of an overall energy imbalance due to poor diet and limited physical activity. Obesity increases the risk for health conditions such as coronary heart disease, type 2 diabetes, cancer, hypertension, dyslipidemia, stroke, liver and gallbladder disease, sleep apnea and respiratory problems, and osteoarthritis.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Physical Inactivity - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Percent of adults reporting no leisure time physical activity, 2008



## CONTEXT

**What It Is:** Physical inactivity is the estimated percent of adults ages 20 and older reporting no leisure time physical activity.

**Where It Comes From:** Estimates of physical inactivity by county were calculated by the CDC's National Center for Chronic Disease Prevention and Health Promotion, Division of Diabetes Translation, using multiple years of Behavioral Risk Factor Surveillance System (BRFSS) data. BRFSS data are representative of the total non-institutionalized U.S. population ages 18 and older living in households with a land-line telephone.


**Importance:** Regular physical activity is one of the most important things one can do for their health. It can help control weight, reduce risk of cardiovascular disease, reduce risk for type 2 diabetes and metabolic syndrome, reduce risk of some cancers, strengthen bones and muscles, improve mental health and mood, improve ability to do daily activities and prevent falls in older adults, and increase chances of living longer (Centers for Disease Control and Prevention, <http://www.cdc.gov/physicalactivity/everyone/health/index.html>).

- Data were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Excessive Drinking - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Percent of adults reporting binge drinking and heavy drinking, 2003-2009



## CONTEXT

**What It Is:** The excessive drinking measure reflects the percent of the adult population that reports either binge drinking, defined as consuming more than 4 (women) or 5 (men) alcoholic beverages on a single occasion in the past 30 days, or heavy drinking, defined as drinking more than 1 (women) or 2 (men) drinks per day on average.

**Where It Comes From:** This measure was calculated by the National Center for Health Statistics using data obtained from the Centers for Disease Control and Prevention’s Behavioral Risk Factor Surveillance System (BRFSS), a random-digit dial survey. BRFSS data are representative of the total non-institutionalized U.S. population ages 18 and older living in households with a land-line telephone. The estimates are based on seven years of data.


**Importance:** Excessive drinking is a risk factor for a number of adverse health outcomes such as alcohol poisoning, hypertension, acute myocardial infarction, sexually transmitted infections, unintended pregnancy, fetal alcohol syndrome, sudden infant death syndrome, suicide, interpersonal violence, and motor vehicle crashes.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Motor Vehicle Crash Death Rate - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Motor vehicle crash deaths per 100,000 population, 2001-2007



## CONTEXT

**What It Is:** Motor vehicle crash deaths are measured as the crude mortality rate per 100,000 population due to on- or off-road accidents involving a motor vehicle. Motor vehicle deaths includes traffic and non-traffic accidents involving motorcycles and 3-wheel motor vehicles; cars; vans; trucks; buses; street cars; ATVs; industrial, agricultural, and construction vehicles; and bikes and pedestrians when colliding with any of the vehicles mentioned. Deaths due to boating accidents and airline crashes are not included in this measure.

**Where It Comes From:** These data were calculated by National Center for Health Statistics (NCHS), part of the Centers for Disease Control and Prevention (CDC), based on data reported to the National Vital Statistics System (NVSS). NCHS used data for a seven-year period to create more robust estimates of cause-specific mortality, particularly for counties with smaller populations.


**Importance:** A strong association has been demonstrated between excessive drinking and alcohol-impaired driving, with approximately 17,000 Americans killed annually in alcohol-related motor vehicle crashes.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Sexually Transmitted Infections - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Number of chlamydia cases (new cases reported) per 100,000 population, 2008



## CONTEXT

**What It Is:** The Sexually Transmitted Infection (STI) rate is measured as chlamydia incidence (the number of new cases reported) per 100,000 population.

**Where It Comes From:** The county-level measures were obtained from the CDC's National Center for Hepatitis, HIV, STD, and TB Prevention.


**Importance:** Chlamydia is the most common bacterial STI in North America and is one of the major causes of tubal infertility, ectopic pregnancy, pelvic inflammatory disease, and chronic pelvic pain. STIs in general are associated with a significantly increased risk of morbidity and mortality, including increased risk of cervical cancer, involuntary infertility, and premature death. However, increases in reported chlamydia infections may reflect the expansion of chlamydia screening, use of increasingly sensitive diagnostic tests, an increased emphasis on case reporting from providers and laboratories, improvements in the information systems for reporting, as well as true increases in disease.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.


Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011

# Teen Birth Rate - A health factor measure focusing on health behaviors

County distribution map for Iowa, Minnesota, Nebraska, North Dakota, and South Dakota



Number of teen births per 1,000 females ages 15 through 19, 2001-2007



## CONTEXT

**What It Is:** Teen births are reported as the number of births per 1,000 female population ages 15 through 19.

**Where It Comes From:** Teen birth rates were obtained from the National Vital Statistics System (NVSS) at the National Center for Health Statistics, part of the Centers for Disease Control and Prevention (CDC).

**Importance:** Teen pregnancy is associated with poor prenatal care and pre-term delivery. Pregnant teens are more likely than older women to receive late or no prenatal care, have gestational hypertension and anemia, and achieve poor maternal weight gain. They are also more likely to have a pre-term delivery and low birth weight, increasing the risk of child developmental delay, illness, and mortality.

- Data and associated context were obtained from County Health Rankings, a key component of the Mobilizing Action Toward Community Health (MATCH) project - a collaboration between the Robert Wood Johnson Foundation and the University of Wisconsin Population Health Institute, <http://www.countyhealthrankings.org/>.

Disclaimer: The data displayed are from the source indicated; we do not vouch for the accuracy of the data or ensure they are the most recent available. The information is intended for personal, non-commercial use. It can be shared freely if it is not used for profit and appropriate acknowledgments are given. This map was prepared by researchers at North Dakota State University in Fargo for the 2011-2013 Fargo-Moorhead Community Health Needs Assessment Collaborative. December 2011